

©American Psychological Association, 2022. This paper is not the copy of record and may not exactly replicate the authoritative document published in the APA journal. The final article is available, upon publication, at: [https://doi.org/ 10.1037/dev0001384](https://doi.org/10.1037/dev0001384)

Connecting Symbolic Fractions to their Underlying Proportions Using Iterative Partitioning

Michelle A. Hurst*, Jacob R. Butts, & Susan C. Levine

Department of Psychology, University of Chicago

Word Count: 9975

Date Accepted: March 16th 2022

* hurstm@uchicago.edu or hurst.mich@gmail.com

Author Note

All materials, data, analysis code, and pre-registration documents are available on the Open Science Framework project page (<https://osf.io/ths6m/>). The study was pre-registered on As Predicted (#19466; <https://aspredicted.org/c7hc6.pdf>). Additionally, we would like to thank the participating schools, programs, and children, including the Chicago Public Schools, for their support of this project.

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

Abstract

Fractions are a challenging mathematics topic for many elementary and middle school students, and even for adults. However, a growing body of developmental research suggests that young children can reason about visually presented proportions, well before fraction instruction, providing insight into how fractions might be introduced to improve learning. We designed a card game to teach 1st and 2nd grade children (N = 195, including a racially and economically diverse sample from the U.S.) about fractions in one of three ways. In the Actively Divided condition we iteratively divided an area model into equal-sized units, in the Pre-Divided condition we used an area model with the end-state of the Actively Divided condition, and in the Non-Divided condition we used a continuous representation of the fraction magnitude that was not divided into unit-sized parts. Children in the Actively Divided condition demonstrated larger improvements matching symbolic fractions and visual fractions (i.e., pie charts) than children in the other two conditions. Post-hoc analyses of children's gameplay revealed that the Actively Divided condition may have provided a more optimal level of difficulty for young children than the Pre-Divided condition, which was particularly difficult, and the Non-Divided condition, which was trivially easy. These differences in gameplay performance provide insights into possible mechanisms for our results. We discuss open research questions highlighted by this work and implications of these findings for both the development of proportional reasoning and fraction learning.

Keywords: proportional reasoning; fractions; card games; non-symbolic fractions; area models

Introduction

Fraction knowledge is an important gatekeeper for success on more advanced math topics, particularly algebra, which in turn is a gatekeeper for further math education (Booth & Newton, 2012; Siegler et al., 2012; Smith, 1996; Spielhagen, 2006). Fractions are also important for the workforce, with over two-thirds of adults in the US using rational numbers for their work (Handel, 2016). Unfortunately, despite the extensive time dedicated to fraction instruction throughout grade school, many students demonstrate a poor understanding of fractions that persists into adulthood (Ciosek & Samborska, 2016; Hecht & Vagi, 2010; National Mathematics Advisory Panel, 2008; Siegler & Lortie-Forgues, 2015; Siegler & Pyke, 2013). Thus, students' poor proficiency with fractions can have lasting consequences on future school success and career trajectories. In contrast to these fraction difficulties, infants and young children have sophisticated intuitions about non-symbolic proportional quantities (Boyer et al., 2008; Boyer & Levine, 2012; Denison et al., 2013; Denison & Xu, 2010; Hurst & Cordes, 2018a; Jeong et al., 2007; Kushnir et al., 2010; McCrink & Wynn, 2007). This contrast between early non-symbolic intuitions and later difficulty during formal fraction education has led researchers to suggest that children's non-symbolic understanding of proportion can, and should, be leveraged to support their symbolic fraction learning (e.g., Boyer & Levine, 2015; Matthews & Hubbard, 2016). However, it is not clear *what kind* of non-symbolic representations should be used or *how* they might support symbolic fraction learning. In the current study we address these questions using a fraction card game that varies the features of visual fraction representations used to help students compare symbolic fraction magnitudes. Importantly, we draw upon prior research investigating the development of children's reasoning with non-symbolic visual proportions to generate clear predictions about the features of non-symbolic area model representations that are likely to best

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

support symbolic fraction learning. Our results have implications for both theories that characterize symbol learning and fraction education, as learning symbolic fractions is particularly difficult.

Understanding how people connect symbols to non-symbolic representations, often referred to as the symbol grounding problem, is a central question of human cognition across many different domains (e.g., Harnad, 1990). In the case of proportion, recent evidence suggests that humans have an analogue non-symbolic representation for proportions and can map that analogue non-symbolic representation to symbols (e.g., Binzak & Hubbard, 2020; Kalra et al., 2020; Lewis et al., 2016). However, understanding how people reason about proportion is further complicated by the variability in how it can be represented, both symbolically (e.g., fractions, decimals) and non-symbolically (e.g., subset of items, part of a whole object). For example, people are better able to compare the magnitudes of decimals than fractions (DeWolf et al., 2014; Hurst & Cordes, 2016, 2018b) and are better able to compare the magnitudes of non-symbolic fractions when the representations are continuous (e.g., line lengths, undivided shapes) rather than discrete (e.g., sets of dots, divided shapes; Boyer et al., 2008; Hurst & Cordes, 2018a; Park et al., 2020). Furthermore, when asked to directly map between symbolic and non-symbolic representations, adults prefer and are better at mapping fractions with discrete rather than continuous representations but are better able at mapping decimals with continuous rather than discrete representations (DeWolf et al., 2015). Together, these findings suggest that although people can map symbolic and non-symbolic representations of proportions to one another, the type of representation impacts this mapping. The current study adds to this growing literature by exploring how specific features of non-symbolic fraction representations, namely how discrete

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

information is presented in an area model, can impact children's ability to learn the meaning of fraction symbols.

In addition to being an important component of theories of proportional reasoning, visual representations have been used in the classroom to support fraction learning for decades and are a recommended component of fraction instruction (Matthews & Hubbard, 2016; National Governors Association Center for Best Practices, 2010; Rau & Matthews, 2017). However, the variability across different representations has resulted in substantial disagreement about the specific type of visual representation that best supports fraction learning. Area models, such as pie charts or rectangles that are divided into units to reflect the denominator and partially colored to reflect the numerator, are likely the most common representation of fraction values. In the United States, partitioning shapes into parts is the first introduction to fractions for many students (National Governors Association Center for Best Practices, 2010). Furthermore, area models are heavily used in US mathematics textbooks (Alajmi, 2012) and both children and pre-service teachers are better able to use area models than other kinds of representations (Kaminski, 2018; Luo et al., 2011). These representations have the benefit of clearly defining the whole (i.e., the entire shape) and conveying the specific part-whole values (i.e., the number of colored units out of the total number of units), but they are limited in their ability to continuously represent improper fractions (i.e., fractions that are larger than one, such as $5/2$) as these numbers necessarily require multiple wholes. As an alternative visual model, researchers and educators have increasingly focused on number lines as a tool for representing numerical magnitudes (e.g., Hamdan & Gunderson, 2017; Siegler et al., 2011; Siegler & Opfer, 2003; Siegler & Ramani, 2009). Some experimental intervention and curriculum studies have found that teaching children to represent fractions on number lines improves their ability to compare fraction magnitudes

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

more so than area models (Gunderson et al., 2019; Hamdan & Gunderson, 2017; Saxe et al., 2013). Furthermore, when 4th to 6th grade children were asked to map a fraction onto a number line, they were significantly less likely to partition or divide up the number line than they were to partition a pie chart (Hurst et al., 2020), suggesting that number lines may be particularly useful for eliciting continuous magnitude-based thinking. However, the question of which visual model is better for learning may not have an all or nothing answer and instead depend heavily on specific features of the model, how that model is used, and what concept we are expecting children to learn (e.g., Ainsworth, 1999; Rau & Matthews, 2017). Thus, in the current study, rather than comparing different types of visual representation (e.g., pie charts vs. number lines), we instead focus on a specific feature of visual representations: the availability and salience of relative units.

Prior work investigating young children's informal reasoning with visual proportions suggests that the availability of discrete countable units does impact their non-symbolic proportional reasoning. Young children who do not yet understand symbolic fractions (e.g., as young as 1st grade) are able to reason about proportional information with continuous area models that are not divided (e.g., 75% red and 25% blue; Hurst & Cordes, 2018a; Jeong et al., 2007). However, when the area model is pre-divided into countable pieces, these same children tend to focus on the number of salient numerator pieces and make systematic errors, such as deciding a spinner with 4/9 red pieces is more likely to land on red than a spinner with 2/3 red pieces because $4 > 2$ even though $4/9 < 2/3$ (e.g., Boyer et al., 2008; Hurst & Cordes, 2018; Jeong et al., 2007). Furthermore, this over-attention to the numerator, at the expense of the relational fraction information, is commonly referred to as the *whole number bias* and continues when children learn symbolic fractions in later grades (e.g., Ni & Zhou, 2005). Taken at face

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

value, this prior work might lead to the hypothesis that continuous undivided area models would also be more effective for symbolic fraction instruction. However, these continuous representations do not convey the specific components of fraction symbols (e.g., both $\frac{3}{4}$ and $\frac{6}{8}$ would have the same continuous representation), and people prefer to map fractions with discrete representations rather than with continuous ones (DeWolf et al., 2015). Thus, although continuous area models may be useful for thinking about proportion magnitude, the utility of these representations for teaching children how to interpret discrete fraction components (such as “ $\frac{3}{4}$ ” being 3 parts out of 4 parts) may be limited.

Thus, given the potential shortcomings of both entirely continuous models, which do not have visible units at all, and entirely discrete models, which highlight the number of units in a way that leads to errors, we investigated a third alternative. We developed an area model that makes discrete countable units available but uses iteration to highlight relative unit size and connect the discrete countable units to continuous magnitude. Prior work has emphasized the importance of building on children’s understanding of measurement, and specifically iterative units, to support fraction learning (e.g., Lamon, 1993; Pitkethly & Hunting, 1996; Sophian, 2007; Tzur & Hunt, 2015), including helping children bridge their understanding of whole numbers and fractions (Boyce & Norton, 2016; Sophian, 2007) and improve their understanding of fraction arithmetic (Braithwaite & Siegler, 2021). Furthermore, the idea of iterating units might be particularly important for helping children move from an immature part-whole conception of fractions to a more advanced measurement conception of fractions (Norton & Wilkins, 2009; Wilkins & Norton, 2018). However, given that children also make systematic errors with the part-whole interpretation of fractions (Miura et al., 1999; Paik & Mix, 2003) and visual proportions (e.g., Boyer et al., 2008), it may be that highlighting relative unit size through

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

iteration can also improve this earlier developing skill of mapping a single symbolic fraction to a part-whole visual proportion. Moreover, this approach allows us to compare a specific feature (the saliency of fraction units) across one type of visual representation (i.e., area models), in order to formulate and test specific theoretical predictions about children's proportional reasoning.

The Current Study

We designed an instructional fraction game, based on the card game War. Although previous studies have shown success in improving fraction learning through modifications of the card game “War” (Gabriel et al., 2012; Leutzinger & Nelson, 1980), we were specifically interested in using the basic game structure to investigate how the availability and salience of units impacts children's fraction reasoning by comparing different area models: entirely continuous, entirely discrete, and a hybrid model that actively highlights the unit-based components of fractions.

We focused on children in 1st and 2nd grade –who have not yet received formal fraction instruction – for three reasons. First, we were interested in measuring different aspects of children's fraction knowledge, including their ability to compare symbolic and visual fractions, their understanding of equivalent fractions, and their more basic knowledge of mapping fractions to part-whole visual representations. There has been a recent surge in research focused on understanding how people reason about fraction magnitudes, both symbolically and non-symbolically, much of which suggests that these skills are critically important components of fraction learning (e.g., Boyer et al., 2008; DeWolf et al., 2014, 2015; DeWolf & Vosniadou, 2015; Hurst & Cordes, 2016, 2018a; Kalra et al., 2020; Möhring et al., 2016; Siegler et al., 2011, 2012). However, even before being able to reason about the relative size of fraction magnitudes,

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

children must learn how to interpret a single fraction, X/Y , as corresponding to X parts out of a whole with Y parts (Norton & Wilkins, 2009; Wilkins & Norton, 2018). The Common Core State Standards describe this skill as a fraction benchmark in 3rd grade, but it is introduced for some simple fractions in 2nd grade (National Governors Association Center for Best Practices, 2010). Thus, we targeted children who were unlikely to have formally been taught symbolic fractions, or at most had been introduced to some simple unit fractions (e.g., “one fourth” or “one half”), so that we could measure their basic understanding of the part-whole interpretation of fractions, as well as their understanding of more advanced fraction concepts, such as relative magnitude and equivalent fractions. Second, prior work has found that 1st and 2nd graders make whole-number based errors when identifying fraction symbols, such as erroneously mapping $2/3$ to an image of 2 black dots and 3 grey dots (equivalent to $2/5$ rather than $2/3$; Miura et al., 1999; Paik & Mix, 2003), but that these errors are malleable (Mix & Paik, 2008). Thus, although iterative units are typically thought to be important for moving beyond the part-whole conception of fractions (e.g., Norton & Wilkins, 2009; Wilkins & Norton, 2018), this age group allows us to investigate whether area models with iterative units are also helpful for preventing whole number based errors when identifying the part-whole interpretation of fraction symbols. Third, and lastly, prior work with non-symbolic proportional reasoning suggests that 6-year-olds, who are typically in 1st grade in the USA, can reason well with visually presented continuous proportions, but struggle with visually presented discrete proportions (e.g., Hurst & Cordes, 2018a; Jeong et al., 2007), making this a critical developmental age group for investigating the utility of visual proportions for scaffolding symbolic fraction learning in a way that aligns with the proportional reasoning literature more broadly.

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

Therefore, in the current manuscript we investigated how training with area models that vary the availability and saliency of discrete units improves children's part-whole interpretation of fractions, measured by their ability to map symbolic fractions onto non-symbolic representations. We predicted that drawing children's attention to the meaning of the numerator and denominator through iterative unit partitioning (Actively Divided condition) would best support children's learning of the mapping between symbolic fractions and visual fractions, in line with other work highlighting the role of partitive unit-based reasoning in early fraction learning (e.g., Boyce & Norton, 2016; Sophian, 2007; Wilkins & Norton, 2018). In contrast, we predicted that continuous, non-partitioned models (Non-Divided condition) would be suboptimal because continuous representations do not convey information about how to connect the visual representation to symbolic fractions and thus may be insufficient for supporting the mapping between these representations. Thus, we predicted that children in the Actively Divided condition would outperform children in the Non-Divided condition. We further predicted that using static and pre-divided models (Pre-Divided condition), which are most similar to those used in fraction instruction, would result in intermediate performance relative to the other conditions as this condition provides the discrete information necessary to make the mapping, but in a way that might lead to whole number-based errors.

In addition to measuring children's ability to map between symbolic and non-symbolic representations of fractions, we also included additional tasks measuring children's ability to compare and equate symbolic and visual fractions. These tasks were included for two reasons: (1) to investigate the generalizability of the benefit of unit iteration for supporting different fraction concepts and (2) to address separate research questions about children's fraction knowledge. Specifically, we were interested in how children's ability to work with fractions

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

differs based on the format of the fraction (symbol vs. visual spatial fraction) and the concept being measured (fraction equivalence vs. magnitude comparison), as well as how these more advanced skills relate to children's ability to map a symbolic fraction to a non-symbolic representation, as a measure of their more basic part-whole knowledge of fractions. We chose to measure these more advanced skills because magnitude comparison is often used as a measure of children's fraction knowledge, allowing us to compare our findings with the broader literature, and equivalent fractions are reliant on the multiplicative relations of numerators and denominators (e.g., $2/4 = 4/8$). However, these additional tasks had low reliability estimates with children performing around chance, suggesting that children in the present study found these tasks very difficult, either because of their immature understanding of fractions and/or general task demands. This is likely due to the young age of these children, which was a deliberate decision based on our developmental hypotheses of when the availability and salience of units is likely to have the largest impact on children's initial learning of fractions. Nevertheless, the age of children in the current study may have undermined our ability to effectively measure these more advanced skills. Therefore, although these tasks and additional research questions were included as part of our pre-registration, we do not report them in the current manuscript. For full transparency a complete write up of the methods and results for these tasks is included in Supplementary Materials and all data is accessible via the OSF project page.

Method

Participants

As preregistered, our final sample included 195 1st and 2nd grade children, randomly assigned to one of three between subject conditions: Non-Divided area model condition ($n = 65$; $M_{\text{age}} = 89.7$ months, range = 77-111 months; 35 1st graders and 30 2nd graders; 21 boys, 44 girls,

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

0 undisclosed/other), Pre-Divided area model condition ($n = 65$; $M_{age} = 90.2$ months, range = 76-107 months (one missing age information); 35 1st graders and 30 2nd graders; 31 boys, 34 girls, 0 undisclosed/other), and Actively Divided area model condition ($n = 65$; $M_{age} = 89.9$ months; range = 77-103 months; 35 1st graders and 30 2nd graders; 31 boys, 34 girls, 0 undisclosed/other). This sample size was chosen a priori (using the *pwr* package, Champely, 2018) to provide 80% power to detect condition differences as small as $f^2 = 0.04$ (considered a *small* effect; Cohen, 1992) in the regression analyses with alpha of 0.05. Four additional children (two 1st graders and two 2nd graders) completed the pre-test but not the post-test (due to absences or refusal), and therefore, in line with our pre-registered plan, are not included in the primary analysis of pre-to-post-test learning but are included in analyses that involve pre-test only. One additional child started pre-test but did not finish the session and is therefore not included in any analyses.

Children were randomly assigned to condition within their grade (1st or 2nd grade; to ensure a similar grade distribution across conditions) and in smaller sets (e.g., a balanced sequence of 12 or 30 assignments at a time) based on the number of children recruited from each testing location and the timing of recruitment and testing across sites. When there was imbalance in condition assignments caused by participant exclusion or experimenter error, those conditions were replaced in subsequent blocks. This process ensured we were maintaining approximate balance across conditions throughout the data collection process and in the overall sample.

Children were recruited from the Chicagoland area from February 2019 to February 2020 through local schools, as well as other community organizations, summer camps, and our university database. We did not collect information about the fraction instruction received by the children for any individual or school. However, based on the Common Core State Standards (National Governors Association Center for Best Practices, 2010), it is unlikely that children had

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

much, if any, formal experience with fraction symbols. As discussed in the introduction, this young age group was targeted in order to investigate the effectiveness of area models with different unit-based features on children's initial knowledge of fraction symbols, prior to formal instruction.

Based on the 84-89% of our sample (depending on the question) that completed the demographic survey¹, 54% of parents reported their children as Black or African American, 31% as White, < 1 % as Native American, <1 % as Asian, 0% as Native Hawaiian or Pacific Islander, 10% as other or more than one race, and 23% identified as Hispanic or Latino/a/x. The sample also came from economically diverse backgrounds, with 50% reporting an annual family income below the Chicago median (which was \$58,247 based on the 2019 US Census; *U.S. Census Bureau QuickFacts*) and 50% reporting the category at or above the Chicago median. More specifically, our sample included: 18% reporting an annual family income of \$15,000 or less, 19% between \$15,000 and \$34,999, 13% between \$35,000 and \$49,999, 7% between \$50,000 and \$74,999, 12% between \$75,000 and \$99,999, and 30% reporting \$100,000 or more. Lastly, the education levels of the parent or guardian who completed the demographic form (84% mothers, 13% fathers, 2% unidentified or nonbinary parent, 1% other guardian) also showed variability: 31% high school or less, 39% associate's or bachelor's degree (or equivalent), and 30% graduate degree.

Parents or legal guardians provided written consent prior to children's participation, and children 7-years-old or older provided formal verbal assent. Parents of children tested in our lab

¹ Individual level data from the demographic survey is not made publicly available because some combinations of information apply to only one individual, making the data identifiable. To provide a general description of the sample, however, the information is presented here for the entire group and separated by condition in the Supplemental Materials Table S6.

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

were given \$10/session as travel compensation and schools and community organizations were given a \$50 gift card for supplies. All participating children were offered a small prize (e.g., sticker or small toy) after each testing session. All testing procedures were approved by the University of Chicago Institutional Review Board under protocol IRB17-1599 “Relational Math Reasoning”.

Design

Children participated in two testing sessions one-on-one with an experimenter, each lasting approximately 30 minutes and occurring approximately seven days apart ($M = 7$ days; Range: 5 days to 10 days; 173 of 195 children had sessions 7 days apart). During session one, children completed a pre-test battery of fraction assessments presented on a laptop computer and then played the fraction card game with the experimenter. In session two, children first played the fraction card game with the experimenter and then completed the post-test battery. The same set of tasks was used at both pre- and post-test across all conditions. These tasks included (in this order): (1) Symbolic-to-Non-Symbolic Mapping task (which of these pictures shows this fraction?), (2) Symbolic Comparison task (which is bigger?), (3) Symbolic Equality task (are these equal or not equal?), (4) Spatial Fraction Comparison task, and (5) Spatial Fraction Equality task. As mentioned in the Introduction, only the Symbolic-to-Non-Symbolic Mapping task will be included in the manuscript and all other tasks are reported in Supplemental because of low reliability estimates (Cronbach’s $\alpha s < 0.53$).

The Fraction Card Game was identical in session one and session two. Children in all three between-subjects conditions completed the same fraction comparisons within the game but the area models differed according to the assigned condition. Given the young age of the target sample and the focus on part-whole area models, all fractions used in the card game and the pre-

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

and post-test measures were proper fractions between 0 and 1 that could be depicted as parts of a single-whole (but potentially presented in unreduced form, e.g., 2/4).

Procedure and Stimuli

Pre-Test Warm Up

Prior to beginning the pre-test, children were asked what they thought a fraction was and their response was written and recorded in summarized form by the experimenter. This task was primarily used as a warm-up question and was not asked at post-test. Children's responses to this pre-test question were double coded to gain better insight into children's fraction knowledge prior to the intervention. The coders classified children's responses as either demonstrating no prior knowledge about fractions (i.e., the child said they did not know or gave an answer that was unrelated to fractions) or as demonstrating some fraction knowledge (i.e., the child's answer reflected at least a partial definition of fractions).

Symbolic to Non-Symbolic Mapping Task (modeled after Miura et al., 1999).

The task was administered at both pre- and post-test and was conducted on a 13-inch laptop computer using PsychoPy2 Version 1.85.4 (Peirce et al., 2019). There were two sets of stimuli, one presented at pre-test and the other presented at post-test, with the set of stimuli assigned to pre- versus post-test counterbalanced across individuals. Each set of stimuli included 8 unique trials with good reliability (Cronbach's $\alpha = .86$ and $.87$ for each set).

On each trial, children were presented with a symbolic fraction centered in the upper half of the screen and four divided pie-charts equally spaced along the bottom half of the screen (see Figure 1A). The fractions (6.5 cm tall, 5.4 cm wide, presented upright with a horizontal bar) only included values between zero and one and the numerators and denominators were single digit

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

integers. The non-symbolic pie charts (4.7 cm diameter) included four, randomly ordered, options, with the numerator pieces colored red and the remaining pieces colored light yellow. On each trial, there were four options: (1) the correct answer (e.g., if the target is $\frac{2}{5}$, then 2 red pieces and 3 yellow pieces; Figure 1A, 3rd option), (2) the numerator + denominator answer (e.g., 2 red pieces and 5 yellow pieces; Figure 1A, 4th option), (3) the correct number of pieces, but the pieces are not equally sized (e.g., 2 red pieces and 3 yellow pieces, but the pieces are different sizes; Figure 1A, 2nd option), and (4) a different fraction with the same denominator (e.g., 4 red pieces and 1 yellow piece; Figure 1A, 1st option).




Figure 1: Example stimuli from the Symbolic to Non-Symbolic Mapping Task (Panel A) and the card game (Panel B), including the “front” of the card (B1), which was identical for all conditions, the “back” of the card in the Actively Divided and Non-Divided conditions (B2), before the experimenter applied the stencil, and the “back” of the card in the Pre-Divided condition (B3).

Children were told they would see a symbolic fraction at the top of the screen and pictures of fractions at the bottom of the screen and their job was to choose which of the pictures showed the fraction at the top. On each trial, children pointed to one of the four options and the experimenter recorded the child’s response by pressing 1, 2, 3, or 4 on the keyboard (corresponding to the position of each option). The trial ended as soon as the experimenter pressed the key. After each trial, a fixation point appeared for 1000ms before the next trial started. Children had as much time as they needed to respond and if they responded that they did

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

not know, they were told to make their best guess. The 8 trials were presented in a random order and children were scored based on proportion correct.

Fraction Card Game

Each card (8.75 cm wide by 5.75 cm tall) had a symbolic fraction on the front and that same symbolic fraction, an equal sign, and a rectangle on the back (see Figure 1B). To distinguish each player's cards, the experimenter's cards had a grey border and the child's cards had a green border. The symbolic fractions (2.5 cm tall by 1.8 cm wide) were presented in upright notation with a horizontal line between the numerator and denominator and included unit fractions and non-unit fractions between zero and one. The rectangle (5 cm wide by 2.5 cm tall) was blank for the Non-Divided and Actively Divided conditions (Figure 1, Panel B2) and was divided into the denominator number of pieces for the Pre-Divided condition (Figure 1, Panel B3).

Children were told they would play a card game with the experimenter where each card had a fraction on it and that on each turn whoever had the card with the bigger fraction would get to keep both cards. At the end of the game whoever had the most cards won. Children were also told that before they decided who would keep the cards, they should put the cards on the designated game board (referred to as the "checking board") and figure out which fraction was bigger. The checking board was a laminated piece of cardstock (15 cm tall by 10 cm wide) with card-sized rectangles on the top and bottom labeled "bigger" and "smaller," respectively. The game consisted of two sections. Section 1 included three trials involving only unit fractions and Section 2 included six trials involving unit and non-unit fractions. Between the two sections there was a transition trial on which children were introduced to non-unit fractions.

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

On each trial, the experimenter pulled a card from their envelope and read the fraction label aloud: “my fraction is [fraction label, e.g., *one fifth*].” The experimenter then helped the child pull a card out of their envelope and read the fraction label aloud: “your fraction is [fraction label, e.g., *one half*].” The cards were prearranged in their respective envelopes so that children would progress through a set order of trials. After both cards were on the table (with only the symbolic fractions visible), the experimenter asked the child to decide which fraction was bigger and to place the card with the bigger fraction in the spot labeled “bigger” and the card with the smaller fraction in the spot labeled “smaller.” The experimenter then recorded the child’s initial response. If the child said they did not know or that the fractions were the same, they were told to make their best guess. After the child’s initial response, the cards were flipped over to reveal the rectangles and the experimenter used a condition-specific approach to fill them in.

In the Non-Divided condition, the experimenter said “my fraction is [fraction label, e.g., *one fifth*], so let’s draw [fraction label, e.g., *one fifth*]” while taking a pre-measured stencil and using a fine-point sharpie to draw a line indicating the fraction. Then, the experimenter said: “now I need to color in this much out of the whole” pointing to the fraction part of the rectangle. Next, the experimenter used a blue colored pencil to color that section and then said: “so that shows [fraction label, e.g., *one fifth*].” This was then repeated for the child’s card. On the child’s card, the experimenter used the stencil and drew the fraction line in the appropriate position (to ensure consistency), but the child colored in their own fraction with the colored pencil.

In the Pre-Divided condition, the experimenter said “my fraction is [fraction label, e.g., *one fifth*]. So my shape is divided into [denominator, e.g., *five*] equal pieces and I need to color in [numerator, e.g., *one*] out of [denominator, e.g., *five*] pieces.” The experimenter then colored in the appropriate number of pieces one at a time with the blue pencil and then said, “so that

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

shows [fraction label, e.g., *one fifth*].” This process was then repeated for the child’s card, but the child colored in their own fraction.

In the Actively Divided condition, the experimenter said “my fraction is [fraction label, e.g., *one fifth*]. So my shape needs to be divided into [denominator, e.g., *five*] equal pieces.” The experimenter then took a pre-made stencil the size of the denominator unit and drew the appropriate number of divisions to fully divide the rectangle (e.g., for $1/5$, divided the rectangle into fifths). After the division the experimenter said: “and I need to color in [numerator, e.g., *one*] out of [denominator, e.g., *five*] pieces” and colored in the appropriate number one at a time with the blue pencil, then saying “so that shows [fraction label, e.g., *one fifth*].” As in the other conditions, this was then repeated for the child’s card, on which the experimenter used the stencil and drew the dividing lines, but the child colored in the fraction.

In all conditions, after both rectangles were filled in, the experimenter then asked: “Now that we can see the fraction pictures, which fraction do you think is bigger?” After the child responded, the experimenter recorded the child’s response and provided verbal feedback (i.e., “that’s right!” if they were correct or “actually, this fraction is bigger”). Lastly, the experimenter said: “that means that [fraction label] is bigger than [fraction label], so [my/your] fraction is biggest and [I/you] get to keep both cards!” and the cards were added to either their own or the child’s pile to count later.

During the transition trial between the two sections, children were asked to think about what would happen if there was more than one piece. The two cards from the previous comparison remained visible and a third card with a “ $1/6$ ” fraction on the front and a removable “3” in the upper corner was introduced, saying “let’s say we had three one-sixths”. The experimenter continued: “when that happens, we can write it like this [move the three to be on

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

top of the one, so it now says “ $3/6$ ”] and we can call it three sixths.” The experimenter then filled in the rectangle and prompted the child to color in the correct amount to show three-sixths using the condition-specific process described earlier. The child was then prompted to compare three-sixths to one-fourth (one of the fractions used in the previous comparison, which was still visible) and the experimenter recorded the child’s response and provided verbal feedback (“you’re right, three sixths is bigger than one fourth” or if the child was incorrect, “actually, three sixths is bigger than one fourth”). Regardless of the child’s accuracy, the child was then prompted to add all three cards (the two from the previous comparison and the new transition card) to their pile.

At the end of the game, the experimenter and child counted the cards they collected (i.e., the ones they won from having the biggest fraction). The game was set up so that the child was always declared the winner, having collected 11 cards (they had the bigger fraction on 5 out of 9 trials + the transition card) versus the experimenter’s 8 cards (they had the bigger fraction on 4 out of 9 trials).

Transparency and Openness

The study design, hypotheses, and analysis plan were pre-registered on aspredicted.org (#19466; <https://aspredicted.org/c7hc6.pdf>). We follow the Journal Article Reporting Standards for Quantitative Designs (Kazak, 2018), including reporting how we determined our sample size, all data exclusions, all manipulations, and all measures (although methodological details and results of some measures and pre-registered research questions are reported only in Supplemental Materials due to low task reliability, as discussed above). Analyses that were not pre-registered are clearly indicated when introduced in the Results section. All data, analysis code, pre-

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

registration documents, and research materials are available on the Open Science Framework (<https://osf.io/th56m/>).

All data processing and analysis was performed in R version 4.0.1 (R Core Team, 2018) using RStudio (version 1.2.5001; R Studio Team, 2016), using packages from the tidyverse (Wickham et al., 2019), as well as *ggpubr* v0.4.0 (Kassambara, 2020a), *rstatix* v0.6.0 (Kassambara, 2020b), *effsize* v0.8.0 (Torchiano, 2018), *psych* v2.0.09 (Revelle, 2020), *apaTables* v2.0.5 (Stanley, 2018), *gt* v0.2.2 (Iannone et al., 2020), and *car* v3.0-9 (Fox & Weisberg, 2019).

Results

Pre-Test Warm-Up Question

Two coders classified children's responses to the initial "What is a fraction?" question as either demonstrating no prior knowledge about fractions (i.e., the child said they did not know or gave an answer that was unrelated to fractions) or as demonstrating some understanding of fractions (i.e., the child's answer reflected at least a partial definition of fractions). The two coders agreed on 97.5% of responses, and a third coder resolved disagreements ($n = 5$). Most children's responses (71.7%) indicated they had no knowledge about fractions prior to our intervention (e.g., "not sure," "subtracting," "food") and the remaining children (28.3%) demonstrated at least some understanding of fractions (e.g., "part of a whole," "pieces of a circle"). Thus, prior to the intervention, the children in our sample demonstrated very limited understanding of what the word "fraction" means.

How Do Different Area Models Impact Children's Understanding of Fraction Symbols?

To address our first research question, and as pre-registered, we used linear regression predicting post-test performance on the symbolic-to-non-symbolic mapping task with pre-test score (continuous covariate), grade (1st vs. 2nd grade), and condition (dummy coded) as

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

predictors. Although not pre-registered, we also compared performance at each time point to chance (0.25) and tested for significant learning in each condition separately using a mixed ANOVA on proportion correct with Session (2: pre and post) as a within-subjects factor and Grade (2: 1st, 2nd) as a between-subjects factor. Pre- and post-test means and standard deviations, as well as the within condition statistical analyses (i.e., one-sample *t*-tests comparing performance to chance and pre- vs. post-test comparisons within condition) are provided in Table 1.

Table 1: Descriptive Statistics in each condition at pre- and post-test

	Pre-Test				Post-Test				Learning
	<i>M</i>	<i>SD</i>	one sample vs .25		<i>M</i>	<i>SD</i>	one sample vs .25		η^2_{partial}
			<i>t</i>	<i>p</i>			<i>t</i>	<i>p</i>	
<i>Symbolic to Non-Symbolic Mapping</i>									
Non-Divided	0.38	0.34	3.07	0.003	0.39	0.36	3.09	0.003	0.002
Pre-Divided	0.40	0.32	3.84	< 0.001	0.41	0.36	3.69	< 0.001	0.002
Actively Divided	0.42	0.36	3.90	< 0.001	0.54	0.37	6.31	< 0.001	0.13*

Note: One-sample *t*-test (*df* = 64) compares children's performance to chance (0.25; 4 options). Although not pre-registered, Learning compares pre vs. post-test performance within each condition, with alpha levels adjusted for multiple comparisons. * *p* < 0.0167, ** *p* < 0.001

The overall regression model was significant, $R^2 = .526$, Adjusted $R^2 = 0.516$, $F(4, 190) = 52.7$, $p < 0.001$, with pre-test score a significant predictor $B = 0.75$, 95% CI [0.65, 0.86], $t(190) = 13.95$, $p < 0.001$, and grade not significant $B = 0.04$, 95% CI [-0.03, 0.12], $t(190) = 1.22$, $p = 0.22$. The critical test is the dummy coded predictors for condition (see Figure 2). When using the Actively Divided condition as the reference group, children in this condition scored significantly higher at post-test than children in the Pre-Divided condition, $B = -0.11$, $SE = 0.04$,

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

95% CI [-0.20, -0.02], $t(190) = -2.41$, $p = 0.02$, and children in the Non-Divided condition, $B = -0.12$, $SE = 0.04$, 95% CI [-0.20, -0.03], $t(190) = -2.57$, $p = 0.01$. Children in the Pre-Divided and Non-Divided conditions (reference = Pre-Divided) did not score significantly differently, $B = -0.01$, $SE = 0.04$, $t(190) = -0.17$, $p = 0.87$.




Figure 2: Performance on the Symbolic to Non-Symbolic Mapping task across condition (x-axis) at pre-test (dark grey, left) and post-test (light grey, right). Plots are violin plots with the mean proportion correct and standard error as error bars. Chance performance (0.25) is indicated with a dotted line.

Next, given our condition differences at post-test, we looked at learning in each of the conditions separately (using Bonferroni adjusted $\alpha = 0.0167$ for the main effect of Session in each of the three conditions). In the Actively Divided condition, there was a main effect of Session, $F(1,63) = 9.26$, $p = 0.003$, $\eta^2_{\text{partial}} = 0.13$, with children scoring significantly higher at post-test than at pre-test. There was also a small main effect of Grade, $F(1,63) = 4.47$, $p = 0.04$, $\eta^2_{\text{partial}} = 0.07$, with 2nd graders, $M = 0.45$, scoring higher than 1st graders, $M = 0.40$. However, there was not a significant interaction between session and grade, $p = 0.24$, $\eta^2_{\text{partial}} = 0.02$. There

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

were not significant main effects of Session in the Pre-Divided condition, $F(1,63) = 0.14$, $p = 0.71$, $\eta^2_{\text{partial}} = 0.002$, or the Non-Divided condition, $F(1,63) = 0.12$, $p = 0.73$, $\eta^2_{\text{partial}} = 0.002$. Neither Grade nor the interaction between Grade and Session were significant in either of these conditions, all $ps > 0.4$, $\eta^2_{\text{partial}} < 0.02$.

To further investigate the change in children's fraction knowledge in the Actively Divided condition, we analyzed the data in two additional ways: change at the individual child level and the pattern of children's errors. Although these analyses are post-hoc and were not pre-registered, they provide additional insight that can generate testable hypotheses for future research.

First, beyond the magnitude of improvement from pre- to post-test, we also calculated the number of children who showed an improvement from pre- to post-test in each condition: 29/65 (45%) in the Actively Divided condition, 21/65 (32%) in the Non-Divided condition, and 24/65 (37%) in the Pre-Divided condition. Although there are numerically more children who improved in the Actively Divided condition than in the other two conditions, this pattern is not statistically significant, $\chi^2(2) = 2.13$, $p = .34$. Thus, the Actively Divided condition may have led to *larger* improvements rather than impacting more children. Consistent with this interpretation, the children who improved in the Actively Divided condition did improve by a larger amount, $M = 0.38$, relative to the Non-Divided condition, $M = 0.23$, $t(44.14) = 2.71$, $p = .01$, and to the Pre-Divided condition, $M = 0.29$, although this comparison was not statistically significant, $t(49.84) = 1.44$, $p = .16$.

Second, the task included carefully selected incorrect responses that correspond to common errors, allowing us to better understand the specific kinds of errors children made and whether children's increase in correct responding corresponded with a decrease in a specific type

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

of error: Unequal Units (correct denominator and numerator, but unequal sized units; option 2 in Figure 1A), Part/Part Interpretation (numerator + denominator, such as 2 red and 3 yellow pieces for 2/5; option 4 in Figure 1A), or Denominator Match (same denominator but different numerator, such as 4 red out of 5 for a target of 2/5; option 1 in Figure 1A). First, the most common error for all three conditions at both pre- and post-test was the Part/Part response (see Figure 3), in line with prior work on children's whole number component bias in this task (Miura et al., 1999; Paik & Mix, 2003). Second, to provide some insight into what aspects of children's fraction knowledge might be shifting in the Actively Divided condition, we analyzed the proportion of trials selecting each type of incorrect response at pre- and post-test. Numerically, the proportion of trials on which children in the Actively Divided condition selected each type of incorrect response decreased from pre- to post-test, but this decrease was only significant for the Unequal Units response, $M_{pre} = .16$, $M_{post} = .10$, $t(64) = -2.31$, $p = .02$, $d = -0.29$, and was not significant for either the Part/Part response, $M_{pre} = .31$, $M_{post} = .29$, $t(64) = -0.71$, $p = .48$, $d = -0.09$, or the Denominator Match response, $M_{pre} = .10$, $M_{post} = .07$, $t(64) = -1.29$, $p = .20$, $d = -0.16$. However, the change from pre- to post-test for the various error types did not significantly differ (all $ps > .40$). Thus, this post-hoc and exploratory analysis reveals that although there is a numerically larger increase in children's attention to equal sized units, the increase in correct responses is not attributable to a decrease in a single type of error, but rather a decrease in errors across all error types.

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS




Figure 3: Mean proportion of trials on the Symbolic to Non-Symbolic Mapping task on which children selected each type of response.

How Did Children’s Performance Vary on the Card Game (Additional RQ)?

To better understand the potential mechanisms supporting children’s learning, we explored differences during the actual game play between conditions. Thus, although not pre-registered, we performed a post-hoc analysis of children’s performance during the card game intervention itself. Specifically, we compared children’s ability to compare symbolic fractions in the three training conditions before and after the condition-specific area models were available.

We isolated analyses to only the first session to assess baseline differences in the use of the three distinct models, before substantial learning could occur. Given that children’s judgements about which fraction was bigger were recorded twice, once when only the symbols were visible (the same across conditions) and again after the area models were constructed for each fraction (which differed by condition), we can investigate how performance on the game

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

differed across conditions as a direct result of the different area models. To see the effect of the specific kind of area model on children's comparison ability, we used an ANOVA with Condition (3: Actively Divided, Pre-Divided, Non-Divided) as a between-subject factor and Judgement Type (2: Symbols Only, With Area Models) as a within-subjects factor on proportion correct (combined across the unit and non-unit fractions within Session 1). Twenty-seven children (out of the 199 who completed Session 1) were excluded from these analyses because they had missing game-play data resulting from the experimenter erroneously not recording at least one of the child's responses during the game. Although the children were excluded for experimenter error, it's possible that the experimenter error was caused by the child's behavior during the game (e.g., more chaotic game play, requiring more behavioral management from the experimenter), resulting in systematic missingness. Notably, both included and excluded children scored similarly on the pre-test symbolic ($M_{\text{included}} = .42$, $M_{\text{excluded}} = .45$) and non-symbolic fraction comparisons ($M_{\text{included}} = .55$, $M_{\text{excluded}} = .64$), the two pre-test tasks most like the game play. Nonetheless, this limitation should be kept in mind when interpreting game play performance from the subset of children with complete game play data. A total of 172 children are included in these analyses: Actively Divided $n = 57$; Pre-Divided $n = 59$; Non-Divided $n = 56$. There was a significant main effect of Condition, $F(2,169) = 19.03$, $p < 0.001$, $\eta^2_{\text{partial}} = 0.18$, Judgement Type, $F(1,169) = 235.13$, $p < 0.001$, $\eta^2_{\text{partial}} = 0.58$, and a Condition by Judgement Type interaction, $F(2,169) = 23.9$, $p < 0.001$, $\eta^2_{\text{partial}} = 0.22$ (see Figure 4).




Figure 4: Performance during the Game Play across the three conditions (x-axis), Comparing performance with symbols only (green, left) vs. with the condition-specific area model (pink, right) during Session 1 only. Plots are violin plots with the mean proportion correct and standard error as error bars. There are not condition differences on symbolic comparisons, but when the areas models are available, there are significant pairwise differences between all conditions.

Given the interaction between Condition and Judgement Type, we conducted two one-way ANOVAs across conditions on the symbolic only judgements and the judgements with area models separately. As would be expected, children's scores did not significantly differ across conditions when making judgements about the fraction symbols alone, $M(SD)_{\text{Actively-Divided}} = 0.50(0.23)$, $M(SD)_{\text{Non-Divided}} = 0.53(0.22)$, $M(SD)_{\text{Pre-Divided}} = 0.49(0.21)$, $F(2,169) = 0.514$, $p = 0.60$, $\eta^2_{\text{partial}} < 0.01$. However, when making judgements with the area models available, there were significant condition differences, $F(2,169) = 48.82$, $p < 0.001$, $\eta^2_{\text{partial}} = 0.37$. In particular, children scored highest in the Non-Divided condition, $M(SD)_{\text{Non-Divided}} = 0.97(0.09)$, followed by the Actively Divided condition, $M(SD)_{\text{Actively-Divided}} = 0.76(0.21)$, and then the Pre-Divided

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

condition, $M(SD)_{\text{Pre-Divided}} = 0.62(0.23)$. All pairwise comparisons were significant using $\alpha = 0.0167$ to adjust for three pair-wise comparisons and Welch's t -tests when there was a significant difference in variance: Non-Divided vs. Actively Divided: $t(74.6) = -6.34, p < 0.001, \text{Cohen's } d = -1.30$; Non-Divided vs. Pre-Divided: $t(74.5) = 10.70, p < 0.001, \text{Cohen's } d = 1.96$; Actively Divided vs. Pre-Divided: $t(114) = 3.31, p = 0.001, \text{Cohen's } d = 0.62$. Thus, replicating work in the proportional reasoning literature more broadly (Boyer et al., 2008; Hurst & Cordes, 2018a), children were more accurate comparing continuous, undivided area models than comparing static, divided area models. However, a novel finding that emerges from this analysis is that comparing models that were actively divided is at an intermediate level, falling between these other two models. We discuss the potential importance of this intermediate difficulty on children's performance in the Discussion.

Discussion

In the current experiment, we report the results of a brief fraction card game that compared the effectiveness of using three different area models that varied in the availability and salience of fraction units to teach 1st and 2nd graders about fractions. As hypothesized, we found that actively divided area models are more effective than strictly discrete or strictly continuous models for improving children's part/whole interpretation of fraction symbols. However, we also hypothesized that discrete area models would result in intermediate performance, between the actively divided and strictly continuous area models. Yet, contrary to our hypothesis, the strictly discrete and strictly continuous area models resulted in similarly low learning. That is, despite the discrete area model providing explicit information about the fraction components and being similar to the kinds of area models commonly used in fraction instruction, playing the game with discrete pre-divided area models did not help children learn the part-whole interpretation of

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

fractions. In fact, the actively divided area model was the only card game intervention to result in significantly higher fraction symbol mapping at post-test. Taken together, our findings align with the hypothesis that highlighting the denominator unit through iteration can support young children's fraction learning, consistent with prior research highlighting the benefits of measurement models for fraction instruction (e.g., Boyce & Norton, 2016; Sophian, 2007). Our findings extend this research by showing that even relative to other discrete part-whole models, actively divided area models support young children's initial part-whole interpretation of symbolic fractions, a pre-cursor skill to understanding relative fraction magnitudes.

Differences in performance during the card game itself provide further insight into children's differential use of area models. During gameplay, children in the Actively Divided condition scored between children in the Non-Divided condition and children in the Pre-Divided condition, highlighting the potential limitations of both entirely continuous and entirely discrete area models. Children who received the continuous area model in the Non-Divided condition were able to complete the card game trivially easily and did not have to confront their errors, and as a result did not receive any feedback or relevant information related to erroneous whole-number strategies. In contrast, children who received the discrete area models in the Pre-Divided condition struggled to compare the fractions, even visually, likely because of their erroneous whole-number based strategies (as has been shown in prior work; Boyer et al., 2008; Ni & Zhou, 2005). Like in the Pre-Divided condition, and unlike in the Non-Divided condition, children in the Actively Divided condition were forced to confront their errors when reasoning about discrete numerical information in the context of fractions (e.g., Ni & Zhou, 2005). Yet, in contrast to the more difficult Pre-Divided condition, the active iteration of units in the Actively Divided condition provided children with the support they needed to confront their errors

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

successfully (e.g., rely on a unit fraction; attend to number of units *and* size of units). Thus, one benefit of the actively divided area model may be that it provided the optimal level of difficulty and scaffolding, in line with desirable difficulty accounts of learning and development (e.g., Bjork, 2018; Schmidt & Bjork, 1992). Together, these findings suggest that the Actively Divided condition provided children with discrete information (rather than omitting it entirely, as in the non-divided area models) in a way that highlighted the size of the denominator units and the relation between unit size, number of units, and fraction size (rather than statically, as in the pre-divided area model), thus preventing children's default attention to countable numerical information only. This pattern of results underscores the importance of considering children's intuitions, namely their whole number-based heuristics and errors, and how best to mitigate them when designing educational materials.

Furthermore, although the importance of measurement models, and iteration in particular, is not new, prior work has primarily focused on the benefits of iteration for more advanced fraction concepts, such as moving beyond part-whole reasoning (e.g., Wilkins & Norton, 2018) and learning fraction arithmetic (e.g., Braithwaite & Siegler, 2021). Here, we extend this prior work to suggest that iteration may also play an important role in supporting even more basic part-whole conceptions of fractions. Taken together then, it may be that the use of measurement models that highlight unit iteration is an important instructional approach for many different aspects of fraction learning, from children's first introduction to part/whole fractions to their later learning of formal fraction arithmetic. However, the current study was limited in its ability to investigate how actively dividing area models might support fraction magnitude knowledge and understanding of equivalent fractions, fraction concepts that are typically taught before fraction arithmetic, because children in the current sample showed low performance on these tasks,

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

resulting in low reliability. Thus, future work is necessary to better understand the role of iteration and units across fraction concepts and for children with different levels of fraction knowledge.

In addition to being a promising approach for supporting children's understanding of symbolic fractions, the current study also extends our understanding of the development of proportional reasoning more generally. Prior work investigating attention to discrete numerical information in both symbolic and non-symbolic proportional reasoning tasks has primarily compared representations that either do or do not have discrete countable information available (e.g., Boyer et al., 2008; DeWolf et al., 2015; Hurst & Cordes, 2016, 2018a). However, the current study suggests that it is not the mere presence of discrete countable information that impacts children's strategy use and/or the effectiveness of the representation as a learning tool. In other words, even though both the pre-divided and actively divided area models provided the same non-symbolic representation of the fraction, they resulted in different patterns of performance and different levels of learning. This suggests that the form of the non-symbolic representation may not be enough to determine how children will interpret them and map them to their symbolic fraction system. Rather, how the form is constructed (in this case, with or without iterative units) may impact children's interpretation of the non-symbolic form itself. Thus, discrete representations can be useful for grounding symbolic fractions, even more useful than continuous representations, but children need support to direct their attention to the relevant information within the discrete representation. Specifically, our study suggests that active iteration of discrete units positively impacts children's attention to relevant denominator information and enhances their ability to accurately map non-symbolic proportions to fraction symbols.

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

This conclusion has implications for education because different instructional mediums offer different opportunities for using representations. For example, an area model presented in a textbook is necessarily a static image, limiting the use of iteration in this context. However, working with manipulatives or informal contexts that involve partitioning and iteration (e.g., equal sharing; Empson, 1999) may lead children to engage with area models in a way that highlights denominator as well as numerator units, while also aligning with children's informal mathematical reasoning. Overall, the current findings highlight the need for work that moves beyond gross level comparisons of discrete vs. continuous area models or area models vs. numbers lines, and instead focuses on the specific features of these representations that are helpful to children's proportional reasoning and symbolic fraction understanding, as well as which features of these representations can lead students astray. In the case of the current study, for example, future research is needed to better understand the mechanisms through which our Actively Divided condition supported learning, including which part(s) of the active division process (e.g., highlighting the denominator, seeing the area model transform, using an iterative division process) is most critical.

Although the actively divided version of the card game training did show success in improving children's symbolic fraction knowledge, there are some limitations, both of the training and for the study as a whole. First, the actively divided condition led to larger gains, but did not impact substantially more children than the other two conditions. Suggesting that, at least in its current form, the training might not be a very robust intervention across many diverse students. However, there are two related limitations of our design that might have contributed to this: our training sessions were short (only two 20- to 30-minute sessions, one week apart) and the children tested were very young. Working with young children was a deliberate decision so

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

that we could investigate children's initial understanding of fractions before they have much (or any) formal fraction input. However, this may have limited our ability to reliably measure and impact more advanced components of children's fraction understanding, such as their ability to compare the magnitude of fractions or judge the equivalence of fractions. The low reliability of these measures and near chance performance of participants further highlight children's difficulty with these tasks. Furthermore, children who were particularly unfamiliar with fractions may have needed more time to benefit from the training. Thus, it may be that additional training and/or training that is threaded throughout the curriculum would have led to larger learning gains and allowed more children to benefit from the intervention. Together, the limited training we provided to children and the difficulty they had with most of the tasks weakened our ability to make conclusions about the effectiveness of our intervention in supporting more advanced skills (although, see Supplemental Materials for details on these tasks) and may have restricted the benefit of the intervention to a subset of children. Additionally, we did not include a delayed post-test in the current study, making it unclear whether the learning gains we found are sustained over time. Future research is necessary to examine whether older children's fraction knowledge may also benefit from our actively divided area model for different and more advanced fraction concepts and whether the extended use of these models would lead to stronger learning gains.

Second, a potentially confounding explanation of our findings is that the Actively Divided training condition may have involved more game play time. Although we did not systematically time the sessions, it is possible that there were systematic condition differences in the time spent with the experimenter because drawing many lines and coloring (Actively Divided condition) likely took longer than either not drawing any lines and only coloring (Pre-Divided

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

condition) or only drawing one line and coloring (Non-Divided condition). However, it is important to note that all sessions were fairly short (as discussed above), meaning that any systematic variation would likely be small. Furthermore, anecdotally, the length of the sessions varied based on many other factors, including individual differences in the child (e.g., how carefully they colored) and the experimenter, which likely overshadowed potential systematic time differences across conditions.

Lastly, the current study carefully compared three different methods of fraction representation but was limited in that only rectangular area models in the training tasks and proper fractions between zero and one were used. It remains an open question whether a similar dynamic and active division would be as effective, or potentially more effective, for other models, such as number lines or pie charts, and when incorporating a wider range of fraction values. One of the potential benefits of number lines over area models is representing fractions beyond one, which require multiple wholes (e.g., $\frac{4}{3}$ requires one complete whole, $\frac{3}{3}$, plus part of a whole, $\frac{1}{3}$). Other work suggests that, in general, number lines are more effective than area models for teaching fraction magnitudes (Gunderson et al., 2019; Hamdan & Gunderson, 2017) and the number line representation may be even more likely to draw on children's measurement concepts (Siegler & Smith, 1997). Thus, it may be that number lines, when paired with the active division that highlights unit-size, may be a particularly effective and generalizable approach for incorporating measurement models into fraction instruction. Furthermore, it is worth noting that the rectangular area model used in the current training differed from typical pie chart representations because the fraction unit was iterated only across one dimension, from left to right, whereas pie charts are typically divided into segments around the center, resulting in units along two dimensions. Although children were able to generalize from the rectangular area

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

models used during the training to circular pie charts used during post-test, whether the training itself would have also been effective with pie charts, or whether the unidirectional iteration is an important feature of these models, is an open question. Importantly, building competency with fractions involves the ability to flexibly and accurately use multiple representations (e.g., Rau & Matthews, 2017). Thus, more research is needed to understand whether these same mechanisms apply to other types of fraction representations and to develop instructional guidelines for how to best integrate the use of multiple representations to teach fractions.

In conclusion, we provided evidence that area models can be effective for children's initial learning of fraction symbols, but that dynamically and iteratively dividing area models is a more effective approach than presenting static divided area models or continuous non-divided area models. Directing children's attention to the relevant features within an area model, in this case through unit iteration, may be necessary for using them effectively. These findings highlight the importance of considering the development of children's intuitions about educationally relevant concepts, including their common incorrect heuristics or systematic errors, prior to formal instruction and how educational materials may interact with those intuitions by directing children's attention to different and more relevant features.

References

- Ainsworth, S. (1999). The functions of multiple representations. *Computers & Education*, 33(2–3), 131–152. [https://doi.org/10.1016/S0360-1315\(99\)00029-9](https://doi.org/10.1016/S0360-1315(99)00029-9)
- Alajmi, A. H. (2012). How do elementary textbooks address fractions? A review of mathematics textbooks in the USA, Japan, and Kuwait. *Educational Studies in Mathematics*, 79(2), 239–261. <https://doi.org/10.1007/s10649-011-9342-1>
- Binzak, J. V., & Hubbard, E. M. (2020). No calculation necessary: Accessing magnitude through decimals and fractions. *Cognition*, 199, 104219. <https://doi.org/10.1016/j.cognition.2020.104219>
- Bjork, R. A. (2018). Being Suspicious of the Sense of Ease and Undeterred by the Sense of Difficulty: Looking Back at Schmidt and Bjork (1992). *Perspectives on Psychological Science*, 13(2), 146–148. <https://doi.org/10.1177/1745691617690642>
- Bonato, M., Fabbri, S., Umiltà, C., & Zorzi, M. (2007). The mental representation of numerical fractions: Real or integer? *Journal of Experimental Psychology: Human Perception and Performance*, 33(6), 1410–1419. <https://doi.org/10.1037/0096-1523.33.6.1410>
- Booth, J. L., & Newton, K. J. (2012). Fractions: Could they really be the gatekeeper's doorman? *Contemporary Educational Psychology*, 37(4), 247–253. <https://doi.org/10.1016/j.cedpsych.2012.07.001>
- Boyce, S., & Norton, A. (2016). Co-construction of fractions schemes and units coordinating structures. *The Journal of Mathematical Behavior*, 41, 10–25. <https://doi.org/10.1016/j.jmathb.2015.11.003>

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

- Boyer, T. W., & Levine, S. C. (2012). Child proportional scaling: Is $1/3=2/6=3/9=4/12$? *Journal of Experimental Child Psychology*, *111*(3), 516–533.
<https://doi.org/10.1016/j.jecp.2011.11.001>
- Boyer, T. W., & Levine, S. C. (2015). Prompting children to reason proportionally: Processing discrete units as continuous amounts. *Developmental Psychology*, *51*(5), 615–620.
<https://doi.org/10.1037/a0039010>
- Boyer, T. W., Levine, S. C., & Huttenlocher, J. (2008). Development of proportional reasoning: Where young children go wrong. *Developmental Psychology*, *44*(5), 1478–1490.
<https://doi.org/10.1037/a0013110>
- Braithwaite, D. W., & Siegler, R. S. (2021). Putting fractions together. *Journal of Educational Psychology*, *113*(3), 556–571. <https://doi.org/10.1037/edu0000477>
- Carpenter, T. P., Corbitt, M. K., Kepner, H. S., Lindquist, M. M., & Reys, R. (1980). RESULTS OF THE SECOND NAEP MATHEMATICS ASSESSMENT: SECONDARY SCHOOL. *The Mathematics Teacher*, *73*(5), 329–338.
- Champely, S. (2018). *pwr: Basic Functions for Power Analysis* (R package version 1.2-2) [Computer software]. <https://CRAN.R-project.org/package=pwr>
- Ciosek, M., & Samborska, M. (2016). A false belief about fractions – What is its source? *The Journal of Mathematical Behavior*, *42*, 20–32.
<https://doi.org/10.1016/j.jmathb.2016.02.001>
- Cohen, J. (1992). A Power Primer. *Psychological Bulletin*, *112*(1), 155.
- Denison, S., Reed, C., & Xu, F. (2013). The emergence of probabilistic reasoning in very young infants: Evidence from 4.5- and 6-month-olds. *Developmental Psychology*, *49*(2), 243–249. <https://doi.org/10.1037/a0028278>

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

- Denison, S., & Xu, F. (2010). Twelve- to 14-month-old infants can predict single-event probability with large set sizes: Twelve- to 14-month-olds can predict single-event probability. *Developmental Science, 13*(5), 798–803. <https://doi.org/10.1111/j.1467-7687.2009.00943.x>
- DeWolf, M., Bassok, M., & Holyoak, K. J. (2015). Conceptual structure and the procedural affordances of rational numbers: Relational reasoning with fractions and decimals. *Journal of Experimental Psychology: General, 144*(1), 127–150. <https://doi.org/10.1037/xge0000034>
- DeWolf, M., Grounds, M. A., Bassok, M., & Holyoak, K. J. (2014). Magnitude comparison with different types of rational numbers. *Journal of Experimental Psychology: Human Perception and Performance, 40*(1), 71–82. <https://doi.org/10.1037/a0032916>
- DeWolf, M., & Vosniadou, S. (2015). The representation of fraction magnitudes and the whole number bias reconsidered. *Learning and Instruction, 37*, 39–49. <https://doi.org/10.1016/j.learninstruc.2014.07.002>
- Empson, S. B. (1999). Equal Sharing and Shared Meaning: The Development of Fraction Concepts in a First-Grade Classroom. *Cognition and Instruction, 17*(3), 283–342.
- Fox, J., & Weisberg, S. (2019). *An {R} Companion to Applied Regression, Third Edition*. [R]. Sage. <https://socialsciences.mcmaster.ca/jfox/Books/Companion/>
- Fazio, L. K., DeWolf, M., & Siegler, R. S. (2016). Strategy use and strategy choice in fraction magnitude comparison. *Journal of Experimental Psychology: Learning, Memory, and Cognition, 42*(1), 1.

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

Gabriel, F., Szucs, D., & Content, A. (2013). The Development of the Mental Representations of the Magnitude of Fractions. *PLoS ONE*, 8(11), e80016.

<https://doi.org/10.1371/journal.pone.0080016>

Gabriel, F., Coché, F., Szucs, D., Carette, V., Rey, B., & Content, A. (2012). Developing Children's Understanding of Fractions: An Intervention Study. *Mind, Brain, and Education*, 6(3), 137–146. <https://doi.org/10.1111/j.1751-228X.2012.01149.x>

Gunderson, E. A., Hamdan, N., Hildebrand, L., & Bartek, V. (2019). Number line unidimensionality is a critical feature for promoting fraction magnitude concepts. *Journal of Experimental Child Psychology*, 187, 104657.

<https://doi.org/10.1016/j.jecp.2019.06.010>

Hamdan, N., & Gunderson, E. A. (2017). The number line is a critical spatial-numerical representation: Evidence from a fraction intervention. *Developmental Psychology*, 53(3), 587–596. <https://doi.org/10.1037/dev0000252>

Handel, M. J. (2016). What do people do at work? *Journal for Labour Market Research*, 49(2), 177–197. <https://doi.org/10.1007/s12651-016-0213-1>

Harnad, S. (1990). The symbol grounding problem. *Physica D: Nonlinear Phenomena*, 42(1–3), 335–346.

Hecht, S. A., & Vagi, K. J. (2010). Sources of Group and Individual Differences in Emerging Fraction Skills. *Journal of Educational Psychology*, 102(4), 843–859.

<https://doi.org/10.1037/a0019824>

Hurst, M. A., Butts, J. R., & Levine, S. C. (2022). Connecting Symbolic Fractions to Continuous Proportion Using a Fraction Card Game. Open Science Framework [Data set].

<https://osf.io/ths6m/>

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

- Hurst, M. A., & Cordes, S. (2016). Rational-number comparison across notation: Fractions, decimals, and whole numbers. *Journal of Experimental Psychology: Human Perception and Performance*, *42*(2), 281–293. <https://doi.org/10.1037/xhp0000140>
- Hurst, M. A., & Cordes, S. (2018a). Attending to relations: Proportional reasoning in 3- to 6-year-old children. *Developmental Psychology*, *54*(3), 428–439. <https://doi.org/10.1037/dev0000440>
- Hurst, M. A., & Cordes, S. (2018b). Children’s understanding of fraction and decimal symbols and the notation-specific relation to pre-algebra ability. *Journal of Experimental Child Psychology*, *168*, 32–48. <https://doi.org/10.1016/j.jecp.2017.12.003>
- Hurst, M. A., Massaro, M., & Cordes, S. (2020). Fraction Magnitude: Mapping Between Symbolic and Spatial Representations of Proportion. *Journal of Numerical Cognition*, in press.
- Hurst, M. A., Relander, C., & Cordes, S. (2016). Biases and Benefits of Number Lines and Pie Charts in Proportion Representation. In A. Papafragou, D. Grodner, D. Mirman, & J. C. Trueswell (Eds.), *Proceedings of the 38th Annual Conference of the Cognitive Science Society* (p. 6). Cognitive Science Society.
- Iannone, R., Cheng, J., & Schloerke, B. (2020). *gt: Easily Create Presentation-Ready Display Tables* (0.2.2) [R]. <https://CRAN.R-project.org/package=gt>
- Jeong, Y., Levine, S. C., & Huttenlocher, J. (2007). The Development of Proportional Reasoning: Effect of Continuous Versus Discrete Quantities. *Journal of Cognition and Development*, *8*(2), 237–256. <https://doi.org/10.1080/15248370701202471>

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

- Kallai, A. Y., & Tzelgov, J. (2009). A generalized fraction: An entity smaller than one on the mental number line. *Journal of Experimental Psychology: Human Perception and Performance*, 35(6), 1845–1864. <https://doi.org/10.1037/a0016892>
- Kalra, P. B., Binzak, J. V., Matthews, P. G., & Hubbard, E. M. (2020). Symbolic fractions elicit an analog magnitude representation in school-age children. *Journal of Experimental Child Psychology*, 195, 104844. <https://doi.org/10.1016/j.jecp.2020.104844>
- Kaminski, J. A. (2018). Effects of visual representations on fraction arithmetic learning. In C. Kalish, M. A. Rau, X. (Jerry) Zhu, & T. T. Rogers (Eds.), *Proceedings of the 40th Annual Meeting of the Cognitive Science Society* (p. 6). https://cognitivesciencesociety.org/wp-content/uploads/2019/01/cogsci18_proceedings.pdf
- Kassambara, A. (2020a). *ggpubr: “ggplot2” Based Publication Ready Plots* (0.2.5) [Computer software]. <https://CRAN.R-project.org/package=ggpubr>
- Kassambara, A. (2020b). *rstatix: Pipe-Friendly Framework for Basic Statistical Tests* (0.4.0) [Computer software]. <https://CRAN.R-project.org/package=rstatix>
- Kazak, A. E. (2018). Editorial: Journal article reporting standards. *American Psychologist*, 73(1), 1–2. <https://doi.org/10.1037/amp0000263>
- Kushnir, T., Xu, F., & Wellman, H. M. (2010). Young Children Use Statistical Sampling to Infer the Preferences of Other People. *Psychological Science*, 21(8), 1134–1140. <https://doi.org/10.1177/0956797610376652>
- Lamon, S. J. (1993). Ratio and Proportion: Connecting Content and Children’s Thinking. *Journal for Research in Mathematics Education*, 24(1), 41. <https://doi.org/10.2307/749385>
- Leutinger, L. P., & Nelson, G. (1980). *Fractions with Models*. 7.

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

- Lewis, M. R., Matthews, P. G., & Hubbard, E. M. (2016). Neurocognitive Architectures and the Nonsymbolic Foundations of Fractions Understanding. In *Development of Mathematical Cognition* (pp. 141–164). Elsevier. <https://doi.org/10.1016/B978-0-12-801871-2.00006-X>
- Luo, F., Lo, J.-J., & Leu, Y.-C. (2011). Fundamental Fraction Knowledge of Preservice Elementary Teachers: A Cross-National Study in the United States and Taiwan: Fundamental Fraction Knowledge. *School Science and Mathematics, 111*(4), 164–177. <https://doi.org/10.1111/j.1949-8594.2011.00074.x>
- Matthews, P. G., & Hubbard, E. M. (2016). Making space for spatial proportions. *Journal of Learning Disabilities, 0022219416679133*.
- McCrink, K., & Wynn, K. (2007). Ratio Abstraction by 6-Month-Old Infants. *Psychological Science, 18*(8), 740–745. <https://doi.org/10.1111/j.1467-9280.2007.01969.x>
- Meert, G., Grégoire, J., & Noël, M.-P. (2010). Comparing the magnitude of two fractions with common components: Which representations are used by 10- and 12-year-olds? *Journal of Experimental Child Psychology, 107*(3), 244–259. <https://doi.org/10.1016/j.jecp.2010.04.008>
- Meert, G., Grégoire, J., Seron, X., & Noël, M.-P. (2012). The mental representation of the magnitude of symbolic and nonsymbolic ratios in adults. *The Quarterly Journal of Experimental Psychology, 65*(4), 702–724. <https://doi.org/10.1080/17470218.2011.632485>
- Miura, I. T., Okamoto, Y., Vlahovic-Stetic, V., Kim, C. C., & Han, J. H. (1999). Language Supports for Children's Understanding of Numerical Fractions: Cross-National Comparisons. *Journal of Experimental Child Psychology, 74*(4), 356–365. <https://doi.org/10.1006/jecp.1999.2519>

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

Mix, K. S., & Paik, J. H. (2008). Do Korean Fraction Names Promote Part-Whole Reasoning?

Journal of Cognition and Development, 9(2), 145–170.

<https://doi.org/10.1080/15248370802022605>

Möhring, W., Newcombe, N. S., Levine, S. C., & Frick, A. (2016). Spatial proportional

reasoning is associated with formal knowledge about fractions. *Journal of Cognition and Development*, 17(1), 67–84.

National Governors Association Center for Best Practices. (2010). *Common core state standards for mathematics*. Retrieved from. Council of Chief State School Officers.

<http://www.corestandards.org/Math/>

National Mathematics Advisory Panel. (2008). *Foundations for success: The final report of the national mathematics advisory panel*. U.S. Department of Education.

Ni, Y., & Zhou, Y.-D. (2005). Teaching and learning fraction and rational numbers: The origins and implications of whole number bias. *Educational Psychologist*, 40(1), 27–52.

Norton, A., & Wilkins, J. L. M. (2009). A quantitative analysis of children's splitting operations and fraction schemes. *The Journal of Mathematical Behavior*, 28(2–3), 150–161.

<https://doi.org/10.1016/j.jmathb.2009.06.002>

Paik, J. H., & Mix, K. S. (2003). US and Korean Children's Comprehension of Fraction Names:

A Reexamination of Cross-National Differences. *Child Development*, 74(1), 144–154.

Park, Y., Viegut, A. A., & Matthews, P. G. (2020). More than the sum of its parts: Exploring the development of ratio magnitude versus simple magnitude perception. *Developmental Science*. <https://doi.org/10.1111/desc.13043>

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

Peirce, J., Gray, J. R., Simpson, S., MacAskill, M., Höchenberger, R., Sogo, H., Kastman, E., &

Lindeløv, J. K. (2019). PsychoPy2: Experiments in behavior made easy. *Behavior*

Research Methods, 51(1), 195–203. <https://doi.org/10.3758/s13428-018-01193-y>

Pitkethly, A., & Hunting, R. (1996). A review of recent research in the area of initial fraction

concepts. *Educational Studies in Mathematics*, 30(1), 5–38.

R Core Team. (2020). *R: A language and environment for statistical computing*. R Foundation

for Statistical Computing. <https://www.R-project.org/>

R Studio Team. (2016). *RStudio: Integrated Development for R*. RStudio Inc.

<http://www.rstudio.com/>

Rau, M. A., & Matthews, P. G. (2017). How to make ‘more’ better? Principles for effective use

of multiple representations to enhance students’ learning about fractions. *ZDM*, 49(4),

531–544. <https://doi.org/10.1007/s11858-017-0846-8>

Revelle, W. (2020). *psych: Procedures for Personality and Psychological Research* (2.0.9)

[Computer software]. <https://CRAN.R-project.org/package=psych>

Saxe, G. B., Diakow, R., & Gearhart, M. (2013). Towards curricular coherence in integers and

fractions: A study of the efficacy of a lesson sequence that uses the number line as the

principal representational context. *ZDM*, 45(3), 343–364. [https://doi.org/10.1007/s11858-](https://doi.org/10.1007/s11858-012-0466-2)

[012-0466-2](https://doi.org/10.1007/s11858-012-0466-2)

Schmidt, R. A., & Bjork, R. A. (1992). New Conceptualizations of Practice: Common Principles

in Three Paradigms Suggest New Concepts for Training. *Psychological Science*, 3(4),

207–218. <https://doi.org/10.1111/j.1467-9280.1992.tb00029.x>

Schneider, M., & Siegler, R. S. (2010). Representations of the magnitudes of fractions. *Journal*

of Experimental Psychology. *Human Perception and Performance*, 36(5), 1227.

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

- Siegel, M., & Smith, J. A. (1997). Toward making representation count in children's conceptions of fractions. *Contemporary Educational Psychology, 22*(1), 1–22.
- Siegler, R. S., Duncan, G. J., Davis-Kean, P. E., Duckworth, K., Claessens, A., Engel, M., Susperreguy, M. I., & Chen, M. (2012). Early Predictors of High School Mathematics Achievement. *Psychological Science, 23*(7), 691–697.
<https://doi.org/10.1177/0956797612440101>
- Siegler, R. S., & Lortie-Forgues, H. (2015). Conceptual knowledge of fraction arithmetic. *Journal of Educational Psychology, 107*(3), 909–918.
<https://doi.org/10.1037/edu0000025>
- Siegler, R. S., & Opfer, J. E. (2003). The Development of Numerical Estimation: Evidence for Multiple Representations of Numerical Quantity. *Psychological Science, 14*(3), 237–243.
- Siegler, R. S., & Pyke, A. A. (2013). Developmental and individual differences in understanding of fractions. *Developmental Psychology, 49*(10), 1994–2004.
<https://doi.org/10.1037/a0031200>
- Siegler, R. S., & Ramani, G. B. (2009). Playing linear number board games—But not circular ones—Improves low-income preschoolers' numerical understanding. *Journal of Educational Psychology, 101*(3), 545–560. <https://doi.org/10.1037/a0014239>
- Siegler, R. S., Thompson, C. A., & Schneider, M. (2011). An integrated theory of whole number and fractions development. *Cognitive Psychology, 62*(4), 273–296.
<https://doi.org/10.1016/j.cogpsych.2011.03.001>
- Smith, J. B. (1996). Does an Extra Year Make Any Difference? The Impact of Early Access to Algebra on Long-Term Gains in Mathematics Attainment. *Educational Evaluation and Policy Analysis, 18*(2), 141–153.

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS

- Sophian, C. (2007). *The origins of mathematical knowledge in childhood*. Routledge. Routledge.
- Spielhagen, F. R. (2006). Closing the Achievement Gap in Math: The Long-Term Effects of Eighth-Grade Algebra. *Journal of Advanced Academics*, 18(1), 34–59.
<https://doi.org/10.4219/jaa-2006-344>
- Spinillo, A. G., & Bryant, P. (1991). Children’s Proportional Judgments: The Importance of “Half.” *Child Development*, 62(3), 427. <https://doi.org/10.2307/1131121>
- Stanley, D. (2018). *apaTables: Create American Psychological Association (APA) Style Tables* (2.0.5) [R]. <https://CRAN.R-project.org/package=apaTables>
- Torchiano, M. (2018). *effsize: Efficient Effect Size Computation*. <https://CRAN.R-project.org/package=effsize>
- Tzur, R., & Hunt, J. (2015). Iteration: Unit Fraction Knowledge and the French Fry Tasks. *Teaching Children Mathematics*, 22(3), 148–157.
<https://doi.org/10.5951/teacchilmath.22.3.0148>
- U.S. Census Bureau *QuickFacts: Chicago city, Illinois*. (n.d.). Retrieved February 11, 2021, from <https://www.census.gov/quickfacts/fact/table/chicagocityillinois/LND110210>
- Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T., Miller, E., Bache, S., Müller, K., Ooms, J., Robinson, D., Seidel, D., Spinu, V., ... Yutani, H. (2019). Welcome to the Tidyverse. *Journal of Open Source Software*, 4(43), 1686.
<https://doi.org/10.21105/joss.01686>
- Wilkins, J. L. M., & Norton, A. (2018). Learning progression toward a measurement concept of fractions. *International Journal of STEM Education*, 5(1), 27.
<https://doi.org/10.1186/s40594-018-0119-2>

CONNECTING SYMBOLIC AND NON-SYMBOLIC FRACTIONS